

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5:

(11) International Publication Number:

WO 92/12151

C07D 453/02, A61K 31/435

A1

(43) International Publication Date:

23 July 1992 (23.07.92)

(21) International Application Number:

PCT/US91/08836

(22) International Filing Date:

4 December 1991 (04.12.91)

(30) Priority data:

639,644

10 January 1991 (10.01.91) US (72) Inventor; and

(75) Inventor/Applicant (for US only): LOWE, John, A., III [US/US]; 28 Coveside Lane, Stonington, New London,

CT 06378 (US).

(74) Agents: RICHARDSON, Peter, C. et al.; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US).

tent), FI, FR (European patent), GB (European patent), GR (European patent), HU, IT (European patent), JP, KR, LU (European patent), MC (European patent), NL (European patent), NO, SE (European patent), US.

(60) Parent Application or Grant

(63) Related by Continuation US

Filed on

639,644 (CON)

(71) Applicant (for all designated States except US): PFIZER INC. [US/US]; 236 East 45nd Street, New York, NY 10017 (US).

(81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (EUROPEAN EUROPEAN EU 10 January 1991 (10.01.91)

Published

With international search report.

(54) Title: N-ALKYL QUINUCLIDINIUM SALTS AS SUBSTANCE P ANTAGONISTS

(57) Abstract

The present invention relates to novel N-alkyl quinuclidinium salts, pharmaceutical compositions comprising such compounds and the use of such compounds in the treatment and prevention inflammatory and gastrointestinal disorders, as well as several other disorders. The N-alkyl quinuclidinium salts of this invention have formula (I), wherein R1, R2, and X are as defined below.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
		FI	Finland	MI.	Mali
AU	Australia .	FR	I-rance	MN	Mongolia
₿B	Bartsalos		• • • • • • • • • • • • • • • • • • • •	· MR	Mauritania
BΕ	Belgium	GA	Gahon	· MW	Malawi
BF	Burkina Faso	GB	United Kingdom		Netherlands
BC	Bulgaria	GN	Guinca	NL	
	•	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil			RO	Romania
CA	Canada	IT	luly	RU	Russian Federation
CF	Central African Republic	JP	Japan		Sudan
CC	Congo	KP	Democratic People's Republic	SD	
	•		of Korua	SE	Sweden
CH	Switzerland	КR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire		•	SU	Soviet Union
CM	Cameroon	Li	Licehtenstein	TD	Chad
cs	Czechoslovakia	LK	Srl Lanka		
DE	Ciernany	LU	Luxembourg	TG	Togo
	•	81C	Monaco	US	United States of America
DK	Denmark				

20

25

30

35

4

:

÷

N-ALKYL QUINUCLIDINIUM SALTS AS SUBSTANCE P ANTAGONISTS

Background of the Invention

The present invention relates to novel N-alkyl quinuclidinium salts, pharmaceutical compositions comprising such compounds and the use of such compounds in the treatment and prevention inflammatory and gastrointestinal disorders, as well as several other disorders. The pharmaceutically active compounds of this invention are substance P antagonists.

Substance P is a naturally occurring undecapeptide belonging to the tachykinin family of peptides, the latter being named because of their prompt stimulatory action on smooth muscle tissue. More specifically, substance P is a pharmacologically active neuropeptide that is produced in mammals (having originally been isolated from gut) and possesses a characteristic amino acid sequence that is illustrated by D. F. Veber et al. in U.S. Patent No. 4,680,283. The wide involvement of substance P and other tachykinins in the pathophysiology of numerous diseases has been amply demonstrated in the art. For instance, substance P has recently been shown to be involved in the transmission of pain or migraine (see B.E.B. Sandberg et al., Journal of Medicinal Chemistry, 25, 1009 (1982)), as well as in central nervous system disorders such as anxiety and schizophrenia, in respiratory and inflammatory diseases such as asthma and rheumatoid arthritis, respectively, in rheumatic diseases such as fibrositis, and in gastrointestinal disorders and diseases of the Gl tract such as ulcerative colitis and Crohn's disease, etc. (see D. Regoli in "Trends in Cluster Headache," edited by F. Sicuteri et al., Elsevier Scientific Publishers, Amsterdam, pp. 85-95 (1987)).

Quinuclidine derivatives and related compounds that exhibit activity as substance P receptor antagonists are referred to in PCT Patent Application PCT/US 89/05338, filed November 20, 1989 and United States Patent Application Serial No. 557,442, filed July 23, 1990, both of which are assigned in common with the present application. Similar compounds are referred to in the PCT patent applications entitled "3-Amino-2-Aryl Quinuclidines" and "Quinuclidine Derivatives" and filed on April 25, 1991 and May 15, 1991, respectively. These applications are also assigned in common with the present application.

Piperidine derivatives and related heterocyclic nitrogen containing compounds that are useful as substance P antagonists are referred to in United States Patent Application Serial No. 619,361, filed November 28, 1990 and United States Patent

25

30

Application Serial No. 590,423, filed September 28, 1990, both of which are assigned in common with the present application.

Summary of the Invention

The present invention relates to compounds of the formula

wherein R^1 is (C_1-C_4) alkyl, aliyi, phenyl- (C_1-C_6) alkyl, HOOC- (C_1-C_{10}) alkyl or (C_1-C_4) alkoxy-OOC- (C_1-C_{10}) alkyl; R^2 is selected from the group consisting of phenyl, thienyl, furyl and pyridyl, each of the foregoing R^2 groups being optionally substituted with from one to three substituents independently selected from the group consisting of cyano, nitro, amino, N-mono- (C_1-C_3) alkylamino, fluorine, chlorine, bromine, trifluoromethyl, (C_1-C_3) alkyl, (C_1-C_3) alkoxy, allyoxy, (C_1-C_3) alkoxy-carbonyl, carboxamido and N,N-di- (C_1-C_3) alkyl-carboxamido; and X is a pharmaceutically acceptable counterion, (e.g., chloride, bromide, fluoride, iodide, mesylate, tosylate or trifluoromethanesulfonate).

Examples of pharmaceutically acceptable counterions are halides (e.g., fluoride, chloride, bromide or iodide), (C₁-C₃)alkyl-mono or di-carboxylates, mesylate, tosylate, arylcarboxylates, (C₁-C₃)alkylsulfonates wherein the alkyl molety may optionally be substituted with one or more fluorine atoms, arylsulfonates, citrate, maleate, fumarate, lactate, malate, sulfates, phosphates, nitrates, tartrate, saccharate and pamoate.

The term "alkyl", as used herein, unless otherwise indicated, includes saturated monovalent hydrocarbon radicals having straight, branched or cyclic moieties or combinations thereof.

Preferred compounds of the formula I are those wherein R² is 2-methoxyphenyl.

The present invention also relates to a pharmaceutical composition for treating or preventing a condition selected from the group consisting of inflammatory diseases (e.g., arthritis, psoriasis, asthma and inflammatory bowel disease), colitis, pain, allergies

WO 92/12151 PCT/US91/08836

such as eczema and rhinitis, chronic obstructive alrways disease, hypersensitivity disorders such as poison ivy, vasospastic diseases such as angina, migraine and Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis, reflex sympathetic dystrophy such as shoulder/hand syndrome, peripheral neuropathy, disorders related to immune enhancement or suppression such as systemic lupus erythematosus, and rheumatic diseases such as fibrositis in a mammal, including a human, comprising an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in treating or preventing such condition, and a pharmaceutically acceptable carrier.

10

25

30

4

The present invention also relates to a method of treating or preventing a condition selected from the group consisting of inflammatory diseases (e.g., arthritis, psoriasis, asthma and inflammatory bowel disease), colitis, pain, allergies such as eczema and rhinitis, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, vasospastic diseases such as angina, migraine and Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis, reflex sympathetic dystrophy such as shoulder/hand syndrome, peripheral neuropathy, disorders related to immune enhancement or suppression such as systemic lupus erythematosus, and rheumatic diseases such as fibrositis in a mammal, including a human, comprising administering to said mammal an amount of a compound of the formula I, or a pharmaceutically acceptable sait thereof, effective in treating or preventing such condition.

The present invention also relates to a pharmaceutical composition for antagonizing the effects of substance P in a mammal, including a human, comprising a substance P antagonizing amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of antagonizing the effects of substance P in a mammal, including a human, comprising administering to said mammal a substance P antagonizing amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof.

The present invention also relates to a pharmaceutical composition for treating or preventing a disorder in a mammal, including a human, resulting from an excess of substance P, comprising a substance P antagonizing amount of a compound of the

30

formula i, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of treating or preventing a disorder in a mammal, including a human, resulting from an excess of substance P, comprising administering to said mammal a substance P antagonizing amount of a compound of the formula I, or a pharmaceutically acceptable sait thereof.

The present invention also relates to a pharmaceutical composition for treating or preventing a condition selected from the group consisting of inflammatory diseases (e.g., arthritis, psoriasis, asthma and inflammatory bowel disease), colitis, pain, allergies such as eczema and rhinitis, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, vasospastic diseases such as angina, migraine and Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis, reflex sympathetic dystrophy such as shoulder/hand syndrome, peripheral neuropathy, disorders related to immune enhancement or suppression such as systemic lupus erythematosus, and rheumatic diseases such as fibrositis in a mammal, including a human, comprising an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in antagonizing the effect of substance P at its receptor site, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of treating or preventing a condition selected from the group consisting of Inflammatory diseases (e.g., arthritis, psoriasis, asthma and inflammatory bowel disease), colitis, pain, allergies such as eczema and rhinitis, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, vasospastic diseases such as angina, migraine and Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis, reflex sympathetic dystrophy such as shoulder/hand syndrome, peripheral neuropathy, disorders related to immune enhancement or suppression such as systemic lupus erythematosus, and rheumatic diseases such as fibrositis in a mammal, including a human, comprising administering to said mammal an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in antagonizing the effect of substance P at its receptor site.

The present invention also relates to a pharmaceutical composition for treating or preventing a disorder in a mammal, including a human, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated

15

20

25

:

neurotransmission, comprising an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in antagonizing the effect of substance P at its receptor site, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of treating or preventing a disorder in mammal, including a human, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising administering to said mammal an amount of a compound of the formula I, or a pharmaceutically acceptable sait thereof, effective in antagonizing the effect of substance P at its receptor site.

The present invention also relates to a pharmaceutical composition for treating or preventing a disorder in a mammal, including a human, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in treating or preventing such disorder, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of treating or preventing a disorder in mammal, including a human, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising administering to said mammal an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in treating or preventing such disorder.

The compounds of the formula I have chiral centers and therefore exist in different enantiomeric forms. This invention relates to all optical isomers and all stereoisomers of compounds of the formula I, and mixtures thereof.

The optically active compounds of formula I are additionally useful as intermediate in the synthesis of the corresponding racemic mixtures and opposite enantiomers.

Formula I above includes compounds identical to those depicted but for the fact that one or more hydrogen or carbon atoms are replaced by radioactive isotopes thereof, (e.g., tritium, carbon-14 or nitrogen-15 isotopes thereof). Such radiolabelled compounds are useful as research and diagnostic tools in metabolism pharmacokinetic studies and in binding assays. Specific applications in research include radioligand binding assays, autoradiography studies and in vivo binding

studies, while specific applications in the diagnostic area include studies of the substance P receptor in humans in <u>in vivo</u> binding in the relevant tissues for inflammation, e.g. immune-type cells or cells that are directly involved in inflammatory bowel disorders and the like.

Detailed Description of the Invention

Compounds of the formula I may be prepared by reacting the corresponding compound of the formula

15

10

5

wherein R² is defined as above, with a compound of the formula R¹X, wherein R₁ is defined as above and X is chloro, fluoro, bromo, iodo, tosyloxy, mesyloxy, or trifluoromethanesulfonyloxy. The reaction is generally carried out in a polar solvent such as ethanol, acetone, dimethylformamide or tetrahydrofuran, at a temperature from about 0°C to about 150°C, preferably at about the reflux temperature of the solvent.

Pressures from about 0.5 atmospheres to about 5 atmospheres are generally acceptable, and ambient pressure, i.e. about 1 atmosphere, is preferred as a matter of convenience.

The novel compounds of the formula I and the pharmaceutically acceptable salts thereof are useful as substance P antagonists, i.e., they possess the ability to antagonize the effects of substance P at its receptor site in mammals, and therefore they are able to function as therapeutic agents in the treatment of the aforementioned disorders and diseases in an afflicted mammal.

The compounds of the formula I which are basic in nature are capable of forming a wide variety of different salts with various inorganic and organic acids. Although such salts must be pharmaceutically acceptable for administration to animals, it is often desirable in practice to initially isolate a compound of the Formula I from the reaction mixture as a pharmaceutically unacceptable salt and then simply convert such

sait to an alternate, pharmaceutically acceptable sait by standard ion exchange methods known to those skilled in the art. In additional, the acid addition saits of the compounds of this invention are readily prepared by treating the appropriate compound of formula I with a substantially equivalent amount of the chosen mineral or organic acid in an aqueous solvent medium or in a suitable organic solvent, such as methanol or ethanol. Upon careful evaporation of the solvent, the desired solid sait is readily obtained.

The compounds of Formula I and their pharmaceutically acceptable salts exhibit substance P receptor-binding activity and therefore are of value in the treatment and prevention of a wide variety of clinical conditions the treatment or prevention of which are effected or facilitated by a decrease in substance P mediated neurotransmission. Such conditions include inflammatory diseases (e.g., arthritis, psoriasis, asthma and inflammatory bowel disease), colitis, pain, allergles such as eczema and rhinitis, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, vasospastic diseases such as angina, migraine and Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis, reflex sympathetic dystrophy such as shoulder/hand syndrome, peripheral neuropathy, disorders related to immune enhancement or suppression such as systemic lupus erythematosus, and rheumatic diseases such as fibrositis. Hence, these compounds are readily adapted to therapeutic use as substance P antagonists for the control and/or treatment of any of the aforesaid clinical conditions in mammals, including humans.

The compounds of the formula I and the pharmaceutically acceptable salts thereof can be administered via either the oral, parenteral or topical routes. In general, these compounds are most desirably administered in dosages ranging from about 5.0 mg up to about 1500 mg per day, although variations will necessarily occur depending upon the weight and condition of the subject being treated and the particular route of administration chosen. However, a dosage level that is in the range of about 0.07 mg to about 21 mg per kg of body weight per day is most desirably employed. Variations may nevertheless occur depending upon the species of animal being treated and its individual response to said medicament, as well as on the type of pharmaceutical formulation chosen and the time period and interval at which such administration is carried out. In some instances, dosage levels below the lower limit of the aforesaid range may be more than adequate, while in other cases still larger doses may be

employed without causing any harmful side effect, provided that such larger doses are first divided into several small doses for administration throughout the day.

c

The compounds of the invention may be administered alone or in combination with pharmaceutically acceptable carriers or diluents by either of the three routes previously indicated, and such administration may be carried out in single or multiple doses. More particularly, the novel therapeutic agents of this invention can be administered in a wide variety of different dosage forms, i.e., they may be combined with various pharmaceutically acceptable inert carriers in the form of tablets, capsules, lozenges, troches, hard candies, powders, sprays, creams, salves, suppositories, jellies, gels, pastes, lotions, ointments, aqueous suspensions, injectable solutions, elixirs, syrups, and the like. Such carriers include solid diluents or fillers, sterile aqueous media and various non-toxic organic solvents, etc. Moreover, oral pharmaceutical compositions can be suitably sweetened and/or flavored. In general, the therapeutically-effective compounds of this invention are present in such dosage forms at concentration levels ranging from about 5.0% to about 70% by weight.

For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dicalcium phosphate and glycine may be employed along with various disintegrants such as starch (and preferably com, potato or tapioca starch), alginic acid and certain complex silicates, together with granulation binders like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tabletting purposes. Solid compositions of a similar type may also be employed as fillers in gelatin capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the active ingredient may be combined with various sweetening or flavoring agents, coloring matter or dyes, and, if so desired, emulsifying and/or suspending agents as well, together with such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

25

30

For parenteral administration, solutions of a therapeutic compound of the present invention in either sesame or peanut oil or in aqueous propylene glycol may be employed. The aqueous solutions should be suitably buffered (preferably pH greater than 8) if necessary and the liquid diluent first rendered isotonic. These aqueous

10

15

20

3

solutions are suitable for intravenous injection purposes. The oily solutions are suitable for intraarticular, intramuscular and subcutaneous injection purposes. The preparation of all these solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well known to those skilled in the art.

Additionally, it is also possible to administer the compounds of the present invention topically when treating inflammatory conditions of the skin and this may preferably be done by way of creams, jellies, gels, pastes, ointments and the like, in accordance with standard pharmaceutical practice.

The activity of the compounds of the present invention as substance P antagonists is determined by their ability to inhibit the binding of substance P at its receptor sites in bovine caudate tissue, employing radioactive ligands to visualize the tachykinin receptors by means of auto-radiography. The substance P antagonizing activity of the herein described compounds may be evaluated by using the standard assay procedure described by M. A. Cascieri et al., as reported in the <u>Journal of Biological Chemistry</u>, Vol. 258, p. 5158 (1983). This method essentially involves determining the concentration of the individual compound required to reduce by 50% the amount of radiolabelled substance P ligands at their receptor sites in said isolated cow tissues, thereby affording characteristic IC₅₀ values for each compound tested.

In this procedure, bovine caudate tissue is removed from a -70°C freezer and homogenized in 50 volumes (w./v.) of an ice-cold 50 mM Tris (i.e., trimethamine which is 2-amino-2-hydroxymethyl-1,3-propanediol) hydrochloride buffer having a pH of 7.7. The homogenate is centrifuged at 30,000 x G for a period of 20 minutes. The pellet is resuspended in 50 volumes of Tris buffer, rehomogenized and then recentrifuged at 30,000 x G for another twenty- minute period. The pellet is then resuspended in 40 volumes of ice-cold 50 mM Tris buffer (pH 7.7) containing 2 mM of calcium chloride, 2 mM of magnesium chloride, 40 g/ml of bacitracin, 4µg/ml of leupeptin, 2µg of chymostatin and 200 g/ml of bovine serum albumin. This step completes the production of the tissue preparation.

The radioligand binding procedure is then carried out in the following manner, viz., by initiating the reaction via the addition of 100 μ l of the test compound made up to a concentration of 1 μ M, followed by the addition of 100 μ l of radioactive ligand made up to a final concentration 0.5 mM and then finally by the addition of 800 μ l of the tissue preparation produced as described above. The final volume is thus 1.0 ml,

20

and the reaction mixture is next vortexed and incubated at room temperature (ca. 20°C) for a period of 20 minutes. The tubes are then filtered using a cell harvester, and the glass fiber filters (Whatman GF/B) are washed four times with 50 mM of Tris buffer (pH 7.7), with the filters having previously been presoaked for a period of two hours prior to the filtering procedure. Radioactivity is then determined in a Beta counter at 53% counting efficiency, and the IC₅₀ values are calculated by using standard statistical methods.

The present invention is illustrated by the following example. It will be understood, however, that the invention is not limited to the specific details of this example.

EXAMPLE 1

(2S,3S)-cis-1-Methyl-2-(diphenylmethyl)-N-((2-methoxyphenyl)methyl)-1-aza-bicyclo[2.2.2]octan-3-amine iodide: To a 50 mL round-bottomed flask equipped with condenser and N₂ inlet were added 500 mg (1.21 mmol) (2S,3S)-cis-2-(diphenylmethyl)-N-((2-methoxyphenyl)methyl)-1-azabicyclo[2.2.2]octan-3-amine and 6 mL ethanol. The solution was heated to near boiling, and 207 mg (1.46 mmol) methyl iodide was added. Heating was continued for 10 min, then the solution was cooled to afford a precipitate, which was filtered and dried. The resulting solid gave mp 239-244°C, 328 mg (49% yield).

 1 H-NMR (δ, CDCl₃): 1.8-2.0 (m, 2H), 2.1-2.3 (m, 2H), 2.47 (s, 3H), 2.59 (m, 1H), 3.04 (dd, J=13,84, 2H) (m, 1H), 3.50 (m, 1H), 3.63 (s, 3H), 3.67 (s, 1H), 3.95 (m, 1H), 4.12 (m, 1H), 4.46 (d, J=11.5, 1H), 5.42 (dd, J=6.5, 11.5, 1H), 6.33, 6.68, and 7.07-7.2 (multiplets, 14H), 7.7-7.9 (broad m, 2H).

¹³C-NMR (δ, CDCl₃): 20.0, 22.4, 23,8, 46.1, 49.4, 53.9, 54.6, 55.3, 55.6, 61.2, 25.71.5, 110.2, 120.4, 126.9, 127.2, 127.9, 128.5, 129.6, 141.7, 143.5, 157.1.

Mass Spec. (%): 426 (1, parent), 259 (31), 245 (46), 142 (45), 121 (100), 91 (62).

25

The title compounds of Examples 2-8 were prepared by a procedure similar to that of Example 1.

EXAMPLE 2

(2S,3S)-cis-1-(4-Carbethoxybutyl)-2-(diphenylmethyl)-N-((2-methoxy-phenyl)methyl)-1-azabicyclo[2.2.2]octan-3-amine iodide:

Prepared in 13% yield, m.p. 85°C.

Mass Spec.: 541 (1, parent), 373 (89), 359 (56), 121 (100), 91 (45).

EXAMPLE 3

(2S,3S)-cis-1-(4-Carboethoxyphenylmethyl)-2-(diphenylmethyl)-N-((2-methoxy-10 phenyl)methyl)-1-azabicycio[2.2.2]octan-3-amine iodide:

Prepared in 13% yield, m.p. 140-145°C.

Anal. Calc'd for C₃₈H₄₃N₂O₃I•1/3H₂O: C 64.40, H 6.21, N 3.95. Found: C 64.05, H 6.16, N 3.88.

EXAMPLE 4

15 (2S,3S)-cis-1-(5-Carbomethoxypentyl)-2-(diphenylmethyl)-N-((2-methoxyphenyl)methyl)-1-azabicyclo[2.2.2]octan-3-amine triflate:

Prepared (using acetonitrile instead of ethanol as solvent) in 5% yield, as an oil. Anal. Calc'd for C₃₆H₄₅N₂O₆SF₃•HCl•1/2H₂O: C 57.32, H 6.55, N 3.71. Found: C 57.29, H 6.48, N 3.68. Found: C 57.29, H 6.48, N 3.68.

EXAMPLE 5

(2S,3S)-cis-1-(5-Carboxypentyl)-2-(diphenylmethyl)-N-((2-methoxyphenyl)methyl)-1-azabicyclo[2.2.2]octan-3-amine triflate:

Prepared by hydrolysis of the above compound with potassium hydroxide in ethanol.

High Res. Mass Spec: Calc'd for C₃₄H₄₃N₂O₃: 527.3278. Found: 527.3268.

EXAMPLE 6

(2S,3S)-cis-1-Allyl-2-(diphenylmethyl)-N-((2-methoxyphenyl)methyl)-1-aza-bicyclo[2.2.2]octan-3-amine bromide:

Prepared in 58% yield, m.p. 150-160°C.

30 Anal. Calc'd for C₃₁H₃₇N₂OBr•1.25H₂O: C 66.96, H 7.16, N 5.04. Found: C 66.95, H 7.06, N 4.97.

EXAMPLE 7

(2S,3S)-cis-1-Benzyl-2-(diphenylmethyl)-N-((2-methoxyphenyl)methyl)-1-aza-bicyclo[2.2.2]octan-3-amine bromide:

Prepared in 53% yield, m.p. 206-208°C.

5 Anal. Calc'd for C₃₅H₃₉N₂OBr•H₂O: C 69.87, H 6.87, N 4.66. Found: C 69.48, H 6.84, N 4.52.

EXAMPLE 8

(2S,3S)-cis-1-(Carboethyoxymethyl)-2-(diphenylmethyl)-N-((2-methoxy-phenyl)methyl)-1-azabicyclo[2.2.2]octan-3-amine bromide:

10 Prepared in 17% yield, m.p. 125-135°C.

Anal. Calc'd for C₃₂H₃₉N₂O₃Br•H₂O: C 64.32, H 6.92, N 4.69. Found: C 64.14, H 6.88, N 4.62.

. 4

CLAIMS

1. A compound of the formula

wherein R^1 is (C_1-C_4) alkyl, allyl, phenyl- (C_1-C_6) alkyl, HOOC- (C_1-C_{10}) alkyl or (C_1-C_4) alkoxy-OOC- (C_1-C_{10}) alkyl; R^2 is selected from the group consisting of phenyl, thienyl, furyl and pyridyl, each of the foregoing R^2 groups being optionally substituted with from one to three substituents independently selected from the group consisting of cyano, nitro, amino, N-mono- (C_1-C_3) alkylamino, fluorine, chlorine, bromine, trifluoromethyl, (C_1-C_3) alkyl, (C_1-C_3) alkoxy, allyoxy, (C_1-C_3) alkoxy-carbonyl, carboxamido and N,N-di- (C_1-C_3) alkyl-carboxamido; and X is a pharmaceutically acceptable counterion, or a pharmaceutically acceptable salt thereof.

- 20 2. A compound according to claim 1 wherein R₂ is 2-methoxyphenyl.
 - 3. A compound according to claim 2 wherein R₁ is methyl.
 - 4. A compound according to claim 3 wherein X is iodide.
- 5. A pharmaceutical composition for treating or preventing a condition selected from the group consisting of inflammatory diseases, arthritis, colitis, pain, allergies, chronic obstructive airways disease, hypersensitivity disorders, vasospastic diseases, fibrosing and collagen diseases, reflex sympathetic dystrophy, peripheral neuropathy, disorders related to immune enhancement or suppression and rheumatic diseases in a mammal, comprising an amount of a compound according to claim 1 effective in preventing or treating such condition and a pharmaceutically acceptable carrier.
 - 6. A method of treating or preventing a condition selected from the group consisting of inflammatory diseases arthritis, colitis, pain, allergies, chronic obstructive airways disease, hypersensitivity disorders, vasospastic diseases, fibrosing and

r

20

25

30

collagen diseases, reflex sympathetic dystrophy, peripheral neuropathy, disorders related to immune enhancement or suppression and rheumatic diseases in a mammal, comprising administering to a mammal in need of such treatment or prevention an amount of a compound according to claim 1 effective in preventing or treating such condition.

- 7. A pharmaceutical composition for antagonizing the effects of substance P in a mammal, comprising a substance P antagonizing effective amount of a compound according to claim 1 and a pharmaceutically acceptable carrier.
- 8. A method of antagonizing the effects of substance P in a mammal, comprising administering to said mammal a substance P antagonizing effective amount of a compound according to claim 1.
- A pharmaceutical composition for treating or preventing a condition in a mammal, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising an amount of a compound according to claim 1 effective in antagonizing the effect of substance P at its receptor site and a pharmaceutically acceptable carrier.
 - 10. A method of treating or preventing a condition in a mammal, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising administering to a mammal in need of such treatment or prevention an amount of a compound according to claim 1, or a pharmaceutically acceptable sait thereof, effective in antagonizing the effect of substance P at its receptor site.
 - 11. A pharmaceutical composition for treating or preventing a condition in a mammal, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, effective in treating or preventing such condition and a pharmaceutically acceptable carrier.
 - 12. A method of treating or preventing a condition in mammal, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising administering to a mammal in need of such treatment or prevention an amount of a compound according to claim 1 effective in treating or preventing such condition.
 - 13. A process for preparing a compound of the formula

wherein R¹ is (C₁-C₄)alkyl, allyl, phenyl-(C₁-C₆)alkyl, HOOC-(C₁-C₁₀)alkyl or (C₁-C₄)alkoxy-OOC-(C₁-C₁₀)alkyl; R² is selected from the group consisting of phenyl, thienyl, furyl and pyridyl, each of the foregoing R² groups being optionally substituted with from one to three substituents independently selected from the group consisting of cyano, nitro, amino, N-mono-(C₁-C₃)alkylamino, fluorine, chlorine, bromine, trifluoromethyl, (C₁-C₃)alkyl, (C₁-C₃)alkoxy, allyoxy, (C₁-C₃)alkoxy-carbonyl, carboxamido and N,N-di-(C₁-C₃)alkyl-carboxamido; and X is a pharmaceutically acceptable counterion;

20 R²

comprising reacting a compound of the formula

25

wherein R^2 is defined as above, with a compound of the formula R^1X , wherein R^1 is defined as above and X is chloro, fluoro, bromo, iodo, tosyloxy, mesyloxy or trifluoromethanesulfonyl.

- 14. A process according to claim 13, wherein the compound of formula I prepared by said process is a compound wherein R² is 2-methoxyphenyl.
 - 15. A process according to claim 13, wherein the compound of formula I prepared by said process is a compound wherein R¹ is methyl.

PCT/US91/08836

-16-

WO 92/12151

- 16. A process according to claim 13 wherein the compound of formula I prepared by said process is a compound wherein X⁻ is iodide.
- 17. A process according to claim 13, wherein the compound of formula I prepared by said process is a compound wherein R¹ is selected from carboethoxybutyl, carboethoxyphenylmethyl, carbomethoxypenyl, carboxypentyl, benzyl, allyl, and carboethoxymethyl.

INTERNATIONAL SEARCH REPORT

International Application No PCT/US 91/08836

I. CLAS	SIFICATIO	N OF SUBJECT MATTER (if several classific	ation symbols apply, indicate all) ⁸	·
According	g to Interna	tional Patent Classification (IPC) or to both Na 453/02, A 61 K 31/435	Monai Classification and IPC	
II. FIELD	S SEARCH	ED Minimum Document	ation Searched 7	
Classificati	ion System		assification Symbols	
TDOS		C 07 D		
IPC5		C 07 D		
		Documentation Searched other to the Extent that such Documents	are included in Fields Searched ⁸	
	MENTS A	ONSIDERED TO BE RELEVANTS		
Category *		ion of Document, ¹¹ with indication, where appr	roprists, of the relevant passages 12	Relevant to Claim No.13
X		1, 9005525 (PFIZER INC.) 31		1-5,7,9,
 ^	se	ee the whole document		11,13-
	1			17
Ì				
Х	WO, A	1, 9005729 (PFIZER INC.) 31	May 1990,	1-5,7,9, 11,13-
	S	ee the whole document		17,13
•				
	1			
A	ED A	2, 0404737 (ISTITUTO DE ANG	SELT S.P.A.)	1-5,7,9,
^		7 December 1990,		11,13-
	S	ee the whole document		17
		000 pag		
	<u> </u>			
		ries of cited documents: 18	"I" later document published after or priority date and not in con- cited to understand the princip	the international filing date lict with the application but
		ining the general state of the art which is not be of particular relevance	INVENTION	
1 "	ing date	ment but published on or after the international	"X" document of particular relevar cannot be considered novel or involve an inventive step	ce, the claimed invention cannot be considered to
"L" do	cument whi nich is cited tation or at	ich may throw doubts on priority claim(s) or i to establish the publication date of another her special resson (as specified)		
0 do		erring to an oral disclosure, use, exhibition or	"Y" document of particular relevant cannot be considered to involve document is combined with on ments, such combination being in the art.	e or more other such docu- g obvious to a person skilled
1		bilished prior to the international filing date but priority data claimed	til tild sir	
IV. CERT	TIFICATION			Penmh Report
Date of th	e Actual Co	empletion of the International Search	Date of Mailing of this international	Sesten valots
1st April 1992 2 4. 04, 92				
Internatio	nal Search	ing Authority	Signature of Authorized Officer	
	EURO	PEAN PATENT OFFICE	(1) 40a	12

Form PCT/ISA/210 (second sheet) (January 1985)

Mme Dagmar FRANK

l noci	DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET) Relevant to Claim No		
ategory *	Citation of Document, with indication, where appropriate, of the relevant passages	KEIEVERT TO CLERKE NO	
	EP, A1, 0159059 (LABORATORI GUIDOTTI S.P.A.) 23 October 1985, see the whole document	1-5,7,9, 11,13- 17	
			
	·		
		ļ	
		1	
		ı.	
	<u> </u>		
		′	
	·		
	·		

FURTHER	INFORMATION CONTINUED FROM THE SECOND SHEET	
		į
	•	
	TOTAL OF STATE POLICE TOTAL TOTAL STATE OF STATE	
	SERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE	
This Intern	ational search report has not been established in respect of certain claims under Article 17(2) (a)	tor the tollowing reasons:
<u> </u>	im numbers. $6.8.\%$ because they relate to subject matter not required to be searched by this Aut	nericht numeral
*10	1,12	
14	thods for treatment of the human or animal body	hy surgery
me or	therapy, as well as diagnostic methods [PCT Ru	le 39.1
(i	v)].	
	2	
2. Cla	im numbers	y with the prescribed y:
•		
	begins they are dependent claims and are not drafted in accordance with the	second and third sen-
3. 1 ten	m numbers	
	·	
VI. 🗌 OE	SERVATIONS WHERE UNITY OF INVENTION IS LACKING ²	
This total	rnational Searching Authority found multiple inventions in this international application as follow	s:
*****	manual osarching Authority tours montple inventions in the international opposition	
i		
1. 🗆 Ata	all required additional search feas were timely paid by the applicant, this international search rep ms of the international application.	ort covers all searchable
i .		
2. ☐ An,	only some of the required additional search fees were timely paid by the applicant, this internation thouse claims of the international application for which fees were paid, specifically claims:	inal search report covers
3. 🔲 👷	required additional search (sees were timely paid by the applicant. Consequently, this international of the invention of the international of the chairs. It is covered by claim numbers:	I search report is restrict-
		•
4. 🗆 🏭	all searchable claims could be searched without effort justifying an additional fee, the internation not invite payment of any additional fee.	al Searching Authority
Remark o		
_	additional search fees were accompanied by applicant's protest. protest accompanied the payment of additional seach fees.	
	historia and the man hat man at an annual and an an annual and an	

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.PCT/US 91/08836

SA 54661

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on The European Patent office is in no way tiable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
WO-A1- 9005525	31/05/90	CA-A- EP-A- WO-A-	2003441 0409931 90/05729	23/05/90 30/01/91 31/05/90
WO-A1- 9005729	31/05/90	CA-A- EP-A- WO-A-	2003441 0409931 90/05525	23/05/90 30/01/91 31/05/90
EP-A2- 0404737	27/12/90	AU-D- CA-A- JP-A-	5764990 2019251 3031279	03/01/91 20/12/90 12/02/91
EP-A1- 0159059	23/10/85	DE-A- JP-A- US-A-	3509460 60209521 4603132	19/09/85 22/10/85 29/07/86

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:
☐ BLACK BORDERS
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
☐ FADED TEXT OR DRAWING
☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
☐ SKEWED/SLANTED IMAGES
☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
GRAY SCALE DOCUMENTS
LINES OR MARKS ON ORIGINAL DOCUMENT
☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
OTHER:

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.